



### □ Prostate biopsy 攝護腺穿刺檢查

1. The patient might experience fever, urinary difficulty, **bloody stool**, **hematuria**, urinary frequency, and other symptoms 3 days after the examination. Daily fluid intake of at least 3000cc is recommended after the examination. Patients with renal disease should follow the fluid intake recommendation by their nephrologist. 手術後三天內可能產生發燒、小便困難、**血便**、**血尿**、頻尿等症狀，請繼續服用藥物，每天飲用至少3000cc開水(有腎疾病者請依醫囑攝取水分)。
2. Please seek medical attention if experiencing the following symptoms: **fever over 38.5 °C**, **trouble urinating**, lower abdominal pain or **persistent bloody stool**. 如發高燒超過38.5°C，無法排尿、下腹痛或**持續血便**的情形，請儘速返院就診。



### □ Vasectomy 輸精管結紮手術

1. There will be slight swelling around the wound after surgery, and ice packing will minimize the swelling. 手術後傷口會有輕微腫脹情形，可冰敷減輕腫脹情形。
2. Please seek medical attention as soon as possible if there is increase hematoma and severe bleeding. 如發現血腫塊和傷口嚴重出血，請儘速返院就診。
3. **After surgery, practice contraception for three months or ejaculate at least 15 times**. Outpatient semen analysis will be arranged to confirm the result of surgery. **手術後仍須避孕三個月或至少射精15次以上**，可至門診檢驗精液，以確認手術成功。
4. Work can be resumed immediately after surgery, but heavy lifting should be avoided within 1-2 days. 手術後可立即恢復工作，但最好在1~2天內避免粗重工作。

### □ Circumcision 包皮環切術

1. There will be slight pain and swelling within 3 days after surgery. The prescribed medication should be taken regularly, and the symptoms usually subside after 3 days. 手術後傷口會有輕微疼痛及瘀腫的情形，應定時服用藥物，通常在三天後，症狀即可消退。
2. Follow the doctor's instructions for routine follow-up and review, and keep the wound clean and dry. It will take 7-10 days for the wound to heal. 傷口應保持清潔乾燥，依照醫師指示，定期回診追蹤複查。傷口約七至十天後即可痊癒。
3. In patients with normal penile erection, the erection might cause pain, swelling and bleeding. In case of minimal bleeding, cold packing and compression will stop the bleeding. If there is persistent bleeding despite above technique, medical attention is needed. 青春期及成年的病患在手術後會因陰莖勃起引起傷口的腫痛出血，若是輕微出血時，只需局部冷敷加壓止血即可，若出血嚴重時，應立即回院就診。

4. Sexual activity can be resumed one month after surgery. 手術一個月後，可以恢復性生活。

### □ Other surgeries 其他手術

1. There is usually pain and swelling after surgery, the symptoms usually resolve in 3 days, and the prescribed medication will help alleviate the symptoms. 手術後傷口會有輕微疼痛及瘀腫的情形，應定時服用藥物，通常在三天後症狀即可消退。
2. The wound will be examined in the scheduled clinic follow-up. It will take 7-10 days for the wound to heal. 傷口在七至十天後即可痊癒。傷口癒合前，應依照醫師指示，定期回診追蹤複查。
3. The wound should always be kept clean and dry. Always replace the wound dressing when the gauze is wet or soaked from discharge. 手術後傷口部分應保持清潔乾燥，請小心不要弄濕紗布，如果紗布被浸濕，可先將紗布拿掉，再換藥即可。
4. Apply ointment and cream according to the doctor's instructions. 傷口請按照醫師指示塗抹藥膏。